

GRADIENT ESTIMATES FOR A NONLINEAR DIFFUSION EQUATION ON COMPLETE MANIFOLDS

JIA-YONG WU

ABSTRACT. Let (M, g) be a complete non-compact Riemannian manifold with the m -dimensional Bakry-Émery Ricci curvature bounded below by a non-positive constant. In this paper, we give a localized Hamilton-type gradient estimate for the positive smooth bounded solutions to the following nonlinear diffusion equation

$$u_t = \Delta u - \nabla \phi \cdot \nabla u - au \log u - bu,$$

where ϕ is a C^2 function, and $a \neq 0$ and b are two real constants. This work generalizes the results of Souplet and Zhang (Bull. London Math. Soc., 38 (2006), pp. 1045-1053) and Wu (Preprint, 2008).

1. INTRODUCTION

Let (M, g) be an n -dimensional non-compact Riemannian manifold with the m -dimensional Bakry-Émery Ricci curvature bounded below. Consider the following diffusion equation:

$$(1.1) \quad u_t = \Delta u - \nabla \phi \cdot \nabla u - au \log u - bu$$

in $B(x_0, R) \times [t_0 - T, t_0] \subset M \times (-\infty, \infty)$, where ϕ is a C^2 function, and $a \neq 0$ and b are two real constants. Eq. (1.1) is closely linked with the gradient Ricci solitons, which are the self-similar solutions to the Ricci flow introduced by Hamilton [3]. Ricci solitons have inspired the entropy and Harnack estimates, the space-time formulation of the Ricci flow, and the reduced distance and reduced volume.

Below we recall the definition of Ricci solitons (see also Chapter 4 of [4]).

Definition 1.1. A Riemannian manifold (M, g) is called a *gradient Ricci soliton* if there exists a smooth function $f : M \rightarrow \mathbb{R}$, sometimes called *potential function*, such that for some constant $c \in \mathbb{R}$, it satisfies

$$(1.2) \quad Ric(g) + \nabla^g \nabla^g f = cg$$

on M , where $Ric(g)$ is the Ricci curvature of manifold M and $\nabla^g \nabla^g f$ is the Hessian of f . A soliton is said to be *shrinking*, *steady* or *expanding* if the constant c is respectively positive, zero or negative.

Suppose that (M, g) be a gradient Ricci soliton, and c, f are described in Definition A. Letting $u = e^f$, under some curvature assumptions, we can derive from

Date: January 1, 2009.

2000 *Mathematics Subject Classification.* Primary 58J35; Secondary 58J35, 58J05. Chinese Library Classification: O175.26; O186.12.

Key words and phrases. local gradient estimate; nonlinear diffusion equation; Bakry-Émery Ricci curvature.

(1.2) that (cf. [5], Eq. (7))

$$(1.3) \quad \Delta u + 2cu \log u = (A_0 - nc)u,$$

for some constant A_0 . Eq. (1.3) is a nonlinear elliptic equation and a special case of Eq. (1.1). For this kind of equations, Ma (see Theorem 1 in [5]) obtained the following result.

Theorem A ([5]). *Let (M, g) be a complete non-compact Riemannian manifold of dimension $n \geq 3$ with Ricci curvature bounded below by the constant $-K := -K(2R)$, where $R > 0$ and $K(2R) \geq 0$, in the metric ball $B_{2R}(p)$. Let u be a positive smooth solution to the elliptic equation*

$$(1.4) \quad \Delta u - au \log u = 0$$

with $a > 0$. Let $f = \log u$ and let $(f, 2f)$ be the maximum among f and $2f$. Then there are two uniform positive constant c_1 and c_2 such that

$$(1.5) \quad \begin{aligned} & |\nabla f|^2 - a(f, 2f) \\ & \leq \frac{n[(n+2)c_1^2 + (n-1)c_1^2(1 + R\sqrt{K}) + c_2]}{R^2} + 2n(|a| + K) \end{aligned}$$

in $B_R(p)$.

Then Yang (see Theorem 1.1 in [6]) extended the above result and obtained the following local gradient estimate for the nonlinear equation (1.1) with $\phi \equiv c_0$, where c_0 is a fixed constant.

Theorem B ([6]). *Let M be an n -dimensional complete non-compact Riemannian Manifold. Suppose the Ricci curvature of M is bounded below by $-K := -K(2R)$, where $R > 0$ and $K(2R) \geq 0$, in the metric ball $B_{2R}(p)$. If u is a positive smooth solution to Eq. (1.1) with $\phi \equiv c_0$ on $M \times [0, \infty)$ and $f = \log u$, then for any $\alpha > 1$ and $0 < \delta < 1$*

$$(1.6) \quad \begin{aligned} & |\nabla f|^2(x, t) - \alpha a f(x, t) - \alpha b - \alpha f_t(x, t) \\ & \leq \frac{n\alpha^2}{2\delta t} + \frac{n\alpha^2}{2\delta} \left\{ \frac{2\epsilon^2}{R^2} + \frac{\nu}{R^2} + \sigma + \frac{\epsilon^2}{R^2}(n-1) \left(1 + R\sqrt{K(2R)} \right) \right. \\ & \quad \left. + \frac{K(2R)}{\alpha-1} + \frac{n\alpha^2\epsilon^2}{8(1-\delta)(\alpha-1)R^2} \right\} \end{aligned}$$

in $B_R(p) \times (0, \infty)$, where $\epsilon > 0$ and $\nu > 0$ are some constants and where $\sigma = a/2$ if $a > 0$; $\sigma = -a$ if $a < 0$.

Recently, the author (see Theorem 1.1 in [2]) used Souplet-Zhang's method in [1] and obtained a localized Hamilton-type gradient estimate for the positive smooth bounded solutions of the equation (1.1) with $\phi \equiv c_0$.

Theorem C ([2]). *Let (M, g) be an n -dimensional non-compact Riemannian manifold with $\text{Ric}(M) \geq -K$ for some constant $K \geq 0$. Suppose that $u(x, t)$ is a positive smooth solution to the parabolic equation (1.1) with $\phi \equiv c_0$ in $Q_{R,T} \equiv B(x_0, R) \times [t_0 - T, t_0] \subset M \times (-\infty, \infty)$. Let $f := \log u$. We also assume that there exists non-negative constants α and δ such that $\alpha - f \geq \delta > 0$. Then there exist*

three dimensional constants $\tilde{c}$, $c(\delta)$ and $c(\alpha, \delta)$ such that

$$(1.7) \quad \frac{|\nabla u|}{u} \leq \left(\frac{\tilde{c}}{R}\beta + \frac{c(\alpha, \delta)}{R} + \frac{c(\delta)}{\sqrt{T}} + c(\delta)(|a| + K)^{1/2} + c(\delta)|a|^{1/2}\beta^{1/2} \right) \left(\alpha - \frac{b}{a} - \log u \right)$$

in $Q_{R/2, T/2}$, where $\beta := \max\{1, |\alpha/\delta - 1|\}$.

The purpose of this paper is to extend Theorem C to the general nonlinear diffusion equation (1.1) via the m -dimensional Bakry-Émery Ricci curvature.

Let us first recall some facts about the m -dimensional Bakry-Émery Ricci curvature (please see [7, 8, 9, 10] for more details). Given an n -dimensional Riemannian manifold (M, g) and a C^2 function ϕ , we may define a symmetric diffusion operator $L := \Delta - \nabla\phi \cdot \nabla$, which is the infinitesimal generator of the Dirichlet form

$$\mathcal{E}(f, g) = \int_M (\nabla f, \nabla g) d\mu, \quad \forall f, g \in C_0^\infty(M),$$

where μ is an invariant measure of L given by $d\mu = e^{-\phi} dx$. It is well-known that L is self-adjoint with respect to the weighted measure $d\mu$.

The ∞ -dimensional Bakry-Émery Ricci curvature $Ric(L)$ is defined by

$$Ric(L) := Ric + Hess(\phi),$$

where Ric and $Hess$ denote the Ricci curvature of the metric g and the Hessian respectively. Following the notation used in [10], we also define the m -dimensional Bakry-Émery Ricci curvature of L on an n -dimensional Riemannian manifold as follows

$$Ric_{m,n}(L) := Ric(L) - \frac{\nabla\phi \otimes \nabla\phi}{m - n},$$

where $m := \dim_{BE}(L)$ is called the Bakry-Émery dimension of L . Note that the number m is not necessarily to be an integer and $m \geq n = \dim M$.

The main result of this paper can be stated in the following:

Theorem 1.2. *Let (M, g) be an n -dimensional non-compact Riemannian manifold with $Ric_{m,n}(L) \geq -K$ for some constant $K \geq 0$. Suppose that $u(x, t)$ is a positive smooth solution to the diffusion equation (1.1) in $Q_{R,T} \equiv B(x_0, R) \times [t_0 - T, t_0] \subset M \times (-\infty, \infty)$. Let $f := \log u$. We also assume that there exists non-negative constants α and δ such that $\alpha - f \geq \delta > 0$. Then there exist three dimensional constants $\tilde{c}$, $c(\delta)$ and $c(\alpha, \delta, m)$ such that*

$$(1.8) \quad \frac{|\nabla u|}{u} \leq \left(\frac{\tilde{c}}{R}\beta + \frac{c(\alpha, \delta, m)}{R} + \frac{c(\delta)}{\sqrt{T}} + c(\delta)(|a| + K)^{1/2} + c(\delta)|a|^{1/2}\beta^{1/2} \right) \left(\alpha - \frac{b}{a} - \log u \right)$$

in $Q_{R/2, T/2}$, where $\beta := \max\{1, |\alpha/\delta - 1|\}$.

We make some remarks on the above theorem below.

Remark 1.3. (i). In Theorem 1.2, it seems that the assumption $\alpha - f \geq \delta > 0$ is reasonable. Because from this assumption, we can get $u \leq e^{\alpha - \delta}$. We say that this upper bound of u can be achieved in some setting. For example, from Corollary 1.2 in [6], we know that positive smooth solutions to the elliptic equation (1.4) with $a < 0$ have $u(x) \leq e^{n/2}$ for all $x \in M$ provided the Ricci curvature of M is non-negative.

(ii). Note that the theorem still holds if m -dimensional Bakry-Émery Ricci curvature is replaced by ∞ -dimensional Bakry-Émery Ricci curvature. In fact this result can be obtained by (2.10) in Section 2.

(iii). Theorem 1.2 generalizes the above mentioned Theorem C. When we choose $\phi \equiv c_0$, we return Theorem C. The proof of our main theorem is based on Souplet-Zhang's gradient estimate and the trick used in [2] with some modifications.

In particular, if $u(x, t) \leq 1$ is a positive smooth solution to the diffusion equation (1.1) with $a < 0$, then we have a simple estimate.

Corollary 1.4. *Let (M, g) be an n -dimensional non-compact Riemannian manifold with $\text{Ric}_{m,n}(L) \geq -K$ for some constant $K \geq 0$. Suppose that $u(x, t) \leq 1$ is a positive smooth solution to the diffusion equation (1.1) with $a < 0$ in $Q_{R,T} \equiv B(x_0, R) \times [t_0 - T, t_0] \subset M \times (-\infty, \infty)$. Then there exist two dimensional constants c and $c(m)$ such that*

$$(1.9) \quad \frac{|\nabla u|}{u} \leq \left(\frac{c(m)}{R} + \frac{c}{\sqrt{T}} + c\sqrt{K + |a|} \right) \left(1 - \frac{b}{a} + \log \frac{1}{u} \right)$$

in $Q_{R/2, T/2}$.

Remark 1.5. We point out that our localized Hamilton-type gradient estimate can be also regarded as the generalization of the result of Souplet-Zhang [1] for the heat equation on complete manifolds. In fact, the above Corollary 1.4 is similar to the result of Souplet-Zhang (see Theorem 1.1 of [1]). From the inequality (4.4) below, we can conclude that if $\phi \equiv c_0$ and $a = 0$, then our result can be reduced to theirs.

The method of proving Theorem 1.2 is the gradient estimate, which is originated by Yau [11] (see also Cheng-Yau [12]), and developed further by Li-Yau [13], Li [14] and Negrin [15]. Then R. S. Hamilton [16] gave an elliptic type gradient estimate for the heat equation. But this type of estimate is a global result which requires the heat equation defined on closed manifolds. Recently, a localized Hamilton-type gradient estimate was proved by Souplet and Zhang [1], which can be viewed as a combination of Li-Yau's Harnack inequality [13] and Hamilton's gradient estimate [16]. In this paper, we obtain a localized Hamilton-type gradient estimate for a general diffusion equation (1.1) as Souplet and Zhang in [1] did for the heat equation on complete manifolds. To prove Theorem 1.2, we mainly follow the arguments of Souplet-Zhang in [1], together with some facts about Bakry-Émery Ricci curvature. Note that the diffusion equation (1.1) is nonlinear. So our case is a little more complicated than theirs.

The structure of this paper is as follows. In Section 2, we will give a basic lemma to prepare for proving Theorem 1.2. Section 3 is devoted to the proof of Theorem 1.2. In Section 4, we will prove Corollary 1.4 in the case $0 < u \leq 1$ with $a < 0$.

2. A BASIC LEMMA

In this section, we will prove the following lemma which is essential in the derivation of the gradient estimate of the equation (1.1). Replacing u by $e^{-b/a}u$, we only need to consider positive smooth solutions of the following diffusion equation:

$$(2.1) \quad u_t = \Delta u - \nabla \phi \cdot \nabla u - au \log u.$$

Suppose that $u(x, t)$ is a positive smooth solution to the diffusion equation (1.1) in $Q_{R,T} \equiv B(x_0, R) \times [t_0 - T, t_0]$. Define a smooth function

$$f(x, t) := \log u(x, t)$$

in $Q_{R,T}$. By (2.1), we have

$$(2.2) \quad \left(L - \frac{\partial}{\partial t} \right) f + |\nabla f|^2 - af = 0.$$

Then we have the following lemma, which is a generalization of the computation carried out in [1, 2].

Lemma 2.1. *Let (M, g) be an n -dimensional non-compact Riemannian manifold with $\text{Ric}_{m,n}(L) \geq -K$ for some constant $K \geq 0$. Let $f(x, t)$ is a smooth function defined on $Q_{R,T}$ satisfying the diffusion equation (2.2). We also assume that there exist non-negative constants α and δ such that $\alpha - f \geq \delta > 0$. Then for all (x, t) in $Q_{R,T}$ the function*

$$(2.3) \quad \omega := |\nabla \log(\alpha - f)|^2 = \frac{|\nabla f|^2}{(\alpha - f)^2}$$

satisfies the following inequality

$$(2.4) \quad \begin{aligned} & \left(L - \frac{\partial}{\partial t} \right) \omega \\ & \geq \frac{2(1 - \alpha) + 2f}{\alpha - f} \langle \nabla f, \nabla \omega \rangle + 2(\alpha - f)\omega^2 + 2(a - K)\omega + \frac{2af}{\alpha - f}\omega. \end{aligned}$$

Proof. By (2.3), we have

$$(2.5) \quad \omega_j = \frac{2f_i f_{ij}}{(\alpha - f)^2} + \frac{2f_i^2 f_j}{(\alpha - f)^3},$$

$$(2.6) \quad \Delta \omega = \frac{2|f_{ij}|^2}{(\alpha - f)^2} + \frac{2f_i f_{ijj}}{(\alpha - f)^2} + \frac{8f_i f_j f_{ij}}{(\alpha - f)^3} + \frac{2f_i^2 f_{jj}}{(\alpha - f)^3} + \frac{6f_i^2 f_j^2}{(\alpha - f)^4}$$

and

$$(2.7) \quad \begin{aligned} L\omega &= \Delta \omega - \phi_j \omega_j \\ &= \frac{2|f_{ij}|^2}{(\alpha - f)^2} + \frac{2f_i f_{ijj}}{(\alpha - f)^2} + \frac{8f_i f_j f_{ij}}{(\alpha - f)^3} + \frac{2f_i^2 f_{jj}}{(\alpha - f)^3} + \frac{6f_i^4}{(\alpha - f)^4} - \frac{2f_{ij} f_i \phi_j}{(\alpha - f)^2} - \frac{2f_i^2 f_j \phi_j}{(\alpha - f)^3} \\ &= \frac{2|f_{ij}|^2}{(\alpha - f)^2} + \frac{2f_i (Lf)_i}{(\alpha - f)^2} + \frac{2(R_{ij} + \phi_{ij}) f_i f_j}{(\alpha - f)^2} + \frac{8f_i f_j f_{ij}}{(\alpha - f)^3} + \frac{2f_i^2 \cdot Lf}{(\alpha - f)^3} + \frac{6f_i^4}{(\alpha - f)^4}, \end{aligned}$$

where $f_i := \nabla_i f$ and $f_{ijj} := \nabla_j \nabla_j \nabla_i f$, etc. By (2.3) and (2.2), we also have

$$(2.8) \quad \begin{aligned} \omega_t &= \frac{2\nabla_i f \cdot \nabla_i [Lf + |\nabla f|^2 - af]}{(\alpha - f)^2} + \frac{2|\nabla f|^2 [Lf + |\nabla f|^2 - af]}{(\alpha - f)^3} \\ &= \frac{2\nabla f \nabla Lf}{(\alpha - f)^2} + \frac{4f_i f_j f_{ij}}{(\alpha - f)^2} - \frac{2a|\nabla f|^2}{(\alpha - f)^2} + \frac{2f_i^2 Lf}{(\alpha - f)^3} + \frac{2|\nabla f|^4}{(\alpha - f)^3} - \frac{2af|\nabla f|^2}{(\alpha - f)^3}. \end{aligned}$$

Combining (2.7) with (2.8), we can get

$$(2.9) \quad \left(L - \frac{\partial}{\partial t}\right) \omega = \frac{2|f_{ij}|^2}{(\alpha - f)^2} + \frac{2(R_{ij} + \phi_{ij})f_i f_j}{(\alpha - f)^2} + \frac{8f_i f_j f_{ij}}{(\alpha - f)^3} + \frac{6f_i^4}{(\alpha - f)^4} \\ - \frac{4f_i f_j f_{ij}}{(\alpha - f)^2} - \frac{2f_i^4}{(\alpha - f)^3} + \frac{2af_i^2}{(\alpha - f)^2} + \frac{2af_i^2}{(\alpha - f)^3}.$$

Noting that $Ric_{m,n}(L) \geq -K$ for some constant $K \geq 0$, we have

$$(2.10) \quad (R_{ij} + \phi_{ij})f_i f_j \geq \frac{|\nabla \phi \cdot \nabla f|^2}{m - n} - K|\nabla f|^2 \geq -K|\nabla f|^2.$$

By (2.5), we have

$$(2.11) \quad \omega_j f_j = \frac{2f_i f_j f_{ij}}{(\alpha - f)^2} + \frac{2f_i^2 f_j^2}{(\alpha - f)^3},$$

and consequently,

$$(2.12) \quad 0 = -2\omega_j f_j + \frac{4f_i f_j f_{ij}}{(\alpha - f)^2} + \frac{4f_i^4}{(\alpha - f)^3},$$

$$(2.13) \quad 0 = \frac{1}{\alpha - f} \left[2\omega_j f_j - \frac{4f_i^4}{(\alpha - f)^3} \right] - \frac{4f_i f_j f_{ij}}{(\alpha - f)^3}.$$

Substituting (2.10) into (2.9) and then adding (2.9) with (2.12) and (2.13), we can get

$$(2.14) \quad \left(L - \frac{\partial}{\partial t}\right) \omega \geq \frac{2|f_{ij}|^2}{(\alpha - f)^2} - \frac{2K|\nabla f|^2}{(\alpha - f)^2} + \frac{4f_i f_j f_{ij}}{(\alpha - f)^3} + \frac{2f_i^4}{(\alpha - f)^4} + \frac{2f_i^4}{(\alpha - f)^3} \\ + \frac{2(1 - \alpha) + 2f}{\alpha - f} f_i \omega_i + \frac{2af_i^2}{(\alpha - f)^2} + \frac{2af_i^2}{(\alpha - f)^3}.$$

Note that $\alpha - f \geq \delta > 0$ implies

$$\frac{2|f_{ij}|^2}{(\alpha - f)^2} + \frac{4f_i f_j f_{ij}}{(\alpha - f)^3} + \frac{2f_i^4}{(\alpha - f)^4} \geq 0.$$

This, together with (2.14), yields the desired estimate (2.4). $\square$

3. PROOF OF THEOREM 1.2

In this section, we will use Lemma 2.1 and the localization technique of Souplet-Zhang [1] to give the elliptic type gradient estimates on the positive and bounded smooth solutions of the diffusion equation (1.1).

Proof. First we give the well-known cut-off function by Li-Yau [13] (see also [1]) as follows. We caution the reader that the calculation is not the same as that in [13] due to the difference of the first-order term.

Let $\psi = \psi(x, t)$ be a smooth cut-off function supported in $Q_{R,T}$ satisfying the following properties:

- (1) $\psi = \psi(d(x, x_0), t) \equiv \psi(r, t)$; $\psi(x, t) = 1$ in $Q_{R/2, T/2}$, $0 \leq \psi \leq 1$;
- (2) ψ is decreasing as a radial function in the spatial variables;
- (3) $\frac{|\partial_r \psi|}{\psi^\epsilon} \leq \frac{C_\epsilon}{R}$, $\frac{|\partial_r^2 \psi|}{\psi^\epsilon} \leq \frac{C_\epsilon}{R^2}$, when $0 < \epsilon < 1$;
- (4) $\frac{|\partial_t \psi|}{\psi^{1/2}} \leq \frac{C}{T}$.

From Lemma 2.1, by a straight forward calculation, we have

$$\begin{aligned}
 & L(\psi\omega) - \frac{2(1-\alpha) + 2f}{\alpha - f} \nabla f \cdot \nabla(\psi\omega) - 2 \frac{\nabla\psi}{\psi} \cdot \nabla(\psi\omega) - (\psi\omega)_t \\
 (3.1) \quad & \geq 2\psi(\alpha - f)\omega^2 - \left[\frac{2(1-\alpha) + 2f}{\alpha - f} \nabla f \cdot \nabla\psi \right] \omega - 2 \frac{|\nabla\psi|^2}{\psi} \omega \\
 & + (L\psi)\omega - \psi_t\omega + 2(a - K)\psi\omega + 2 \frac{af}{\alpha - f} \psi\omega.
 \end{aligned}$$

Let (x_1, t_1) be a point where $\psi\omega$ achieves the maximum. By Li-Yau [13], without loss of generality we assume that x_1 is not in the cut-locus of M . Then at this point, we have

$$L(\psi\omega) \leq 0, \quad (\psi\omega)_t \geq 0, \quad \nabla(\psi\omega) = 0.$$

Hence at (x_1, t_1) , by (3.1), we get

$$\begin{aligned}
 (3.2) \quad 2\psi(\alpha - f)\omega^2(x_1, t_1) & \leq \left\{ \left[\frac{2(1-\alpha) + 2f}{\alpha - f} \nabla f \cdot \nabla\psi \right] \omega + 2 \frac{|\nabla\psi|^2}{\psi} \omega - (L\psi)\omega \right. \\
 & \left. + \psi_t\omega - 2(a - K)\psi\omega - 2 \frac{af}{\alpha - f} \psi\omega \right\} (x_1, t_1).
 \end{aligned}$$

In the following, we will introduce the upper bounds for each term of the right-hand side (RHS) of (3.2). Following similar arguments of Souplet-Zhang ([1], pp. 1050-1051), we have the estimates of the first term of the BHS of (3.2)

$$\begin{aligned}
 (3.3) \quad & \left[\frac{2f}{\alpha - f} \nabla f \cdot \nabla\psi \right] \omega \\
 & \leq 2|f| \cdot |\nabla\psi| \cdot \omega^{3/2} = 2[\psi(\alpha - f)\omega^2]^{3/4} \cdot \frac{|f| \cdot |\nabla\psi|}{[\psi(\alpha - f)]^{3/4}} \\
 & \leq \psi(\alpha - f)\omega^2 + \tilde{c} \frac{(f|\nabla\psi|)^4}{[\psi(\alpha - f)]^3} \leq \psi(\alpha - f)\omega^2 + \tilde{c} \frac{f^4}{R^4(\alpha - f)^3}
 \end{aligned}$$

and

$$\begin{aligned}
 (3.4) \quad & \left[\frac{2(1-\alpha)}{\alpha - f} \nabla f \cdot \nabla\psi \right] \omega \\
 & \leq 2|1 - \alpha| |\nabla\psi| \omega^{3/2} = (\psi\omega^2)^{3/4} \cdot \frac{2|1 - \alpha| |\nabla\psi|}{\psi^{3/4}} \\
 & \leq \frac{\delta}{12} \psi\omega^2 + c(\alpha, \delta) \left(\frac{|\nabla\psi|}{\psi^{3/4}} \right)^4 \leq \frac{\delta}{12} \psi\omega^2 + \frac{c(\alpha, \delta)}{R^4}.
 \end{aligned}$$

For the second term of the RHS of (3.2), we have

$$\begin{aligned}
 (3.5) \quad & 2 \frac{|\nabla\psi|^2}{\psi} \omega = 2\psi^{1/2}\omega \cdot \frac{|\nabla\psi|^2}{\psi^{3/2}} \leq \frac{\delta}{12} \psi\omega^2 + c(\delta) \left(\frac{|\nabla\psi|^2}{\psi^{3/2}} \right)^2 \\
 & \leq \frac{\delta}{12} \psi\omega^2 + \frac{c(\delta)}{R^4}.
 \end{aligned}$$

For the third term of the RHS of (3.2), since $\text{Ric}_{m,n}(L) \geq -K$, by the generalized Laplacian comparison theorem (see [9] or [10]),

$$Lr \leq (m - 1)\sqrt{K} \coth(\sqrt{K}r).$$

Consequently, we have

$$\begin{aligned}
(3.6) \quad & -(L\psi)\omega = -\left[(\partial_r\psi)Lr + (\partial_r^2\psi) \cdot |\nabla r|^2\right]\omega \\
& \leq -\left[\partial_r\psi(m-1)\sqrt{K}\coth(\sqrt{K}r) + \partial_r^2\psi\right]\omega \\
& \leq -\left[\partial_r\psi(m-1)\left(\frac{1}{r} + \sqrt{K}\right) + \partial_r^2\psi\right]\omega \\
& \leq \left[|\partial_r^2\psi| + 2(m-1)\frac{|\partial_r\psi|}{R} + (m-1)\sqrt{K}|\partial_r\psi|\right]\omega \\
& \leq \psi^{1/2}\omega\frac{|\partial_r^2\psi|}{\psi^{1/2}} + \psi^{1/2}\omega 2(m-1)\frac{|\partial_r\psi|}{R\psi^{1/2}} + \psi^{1/2}\omega(m-1)\frac{\sqrt{K}|\partial_r\psi|}{\psi^{1/2}} \\
& \leq \frac{\delta}{12}\psi\omega^2 + c(\delta, m)\left[\left(\frac{|\partial_r^2\psi|}{\psi^{1/2}}\right)^2 + \left(\frac{|\partial_r\psi|}{R\psi^{1/2}}\right)^2 + \left(\frac{\sqrt{K}|\partial_r\psi|}{\psi^{1/2}}\right)^2\right] \\
& \leq \frac{\delta}{12}\psi\omega^2 + \frac{c(\delta, m)}{R^4} + \frac{c(\delta, m)K}{R^2}.
\end{aligned}$$

Now we estimate the fourth term:

$$\begin{aligned}
(3.7) \quad & |\psi_t|\omega = \psi^{1/2}\omega\frac{|\psi_t|}{\psi^{1/2}} \leq \frac{\delta}{12}(\psi^{1/2}\omega)^2 + c(\delta)\left(\frac{|\psi_t|}{\psi^{1/2}}\right)^2 \\
& \leq \frac{\delta}{12}\psi\omega^2 + \frac{c(\delta)}{T^2}.
\end{aligned}$$

Notice that we have used Young's inequality below in obtaining (3.3)-(3.7):

$$ab \leq \frac{a^p}{p} + \frac{b^q}{q}, \quad \forall p, q > 0 \text{ with } \frac{1}{p} + \frac{1}{q} = 1.$$

Finally, we estimate the last two terms:

$$(3.8) \quad -2(a-K)\psi\omega \leq 2(|a|+K)\psi\omega \leq \frac{\delta}{12}\psi\omega^2 + c(\delta)(|a|+K)^2;$$

and

$$(3.9) \quad -2\frac{af}{\alpha-f}\psi\omega \leq 2\frac{|a|\cdot|f|}{\alpha-f}\psi\omega \leq \frac{\delta}{12}\psi\omega^2 + c(\delta)a^2\frac{f^2}{(\alpha-f)^2}.$$

Substituting (3.3)-(3.9) to the RHS of (3.2) at (x_1, t_1) , we get

$$\begin{aligned}
(3.10) \quad & 2\psi(\alpha-f)\omega^2 \leq \psi(\alpha-f)\omega^2 + \frac{\tilde{c}f^4}{R^4(\alpha-f)^3} + \frac{\delta}{2}\psi\omega^2 + \frac{c(\alpha, \delta)}{R^4} + \frac{c(\delta)}{R^4} + \frac{c(\delta, m)}{R^4} \\
& + \frac{c(\delta, m)K}{R^2} + \frac{c(\delta)}{T^2} + c(\delta)(|a|+K)^2 + c(\delta)a^2\frac{f^2}{(\alpha-f)^2}.
\end{aligned}$$

Recall that $\alpha-f \geq \delta > 0$, (3.10) implies

$$\begin{aligned}
(3.11) \quad & \psi\omega^2(x_1, t_1) \leq \tilde{c}\frac{f^4}{R^4(\alpha-f)^4} + \frac{1}{2}\psi\omega^2(x_1, t_1) + \frac{c(\alpha, \delta)}{R^4} + \frac{c(\delta, m)}{R^4} + \frac{c(\delta, m)K}{R^2} \\
& + \frac{c(\delta)}{T^2} + c(\delta)(|a|+K)^2 + c(\delta)a^2\frac{f^2}{(\alpha-f)^2}.
\end{aligned}$$

Furthermore, we need to estimate the RHS of (3.11). If $f \leq 0$ and $\alpha \geq 0$, then we have

$$(3.12) \quad \frac{f^4}{(\alpha - f)^4} \leq 1, \quad \frac{f^2}{(\alpha - f)^2} \leq 1;$$

if $f > 0$, by the assumption $\alpha - f \geq \delta > 0$, we know that

$$(3.13) \quad \frac{f^4}{(\alpha - f)^4} \leq \frac{(\alpha - \delta)^4}{\delta^4} = \left(\frac{\alpha}{\delta} - 1\right)^4, \quad \frac{f^2}{(\alpha - f)^2} \leq \left(\frac{\alpha}{\delta} - 1\right)^2.$$

Plugging (3.12) (or (3.13)) into (3.11), we obtain

$$(3.14) \quad (\psi\omega^2)(x_1, t_1) \leq \frac{\tilde{c}\beta^4 + c(\alpha, \delta, m)}{R^4} + \frac{c(\delta, m)K}{R^2} + \frac{c(\delta)}{T^2} + c(\delta)(|a| + K)^2 + c(\delta)a^2\beta^2,$$

where $\beta := \max\{1, |\alpha/\delta - 1|\}$.

The above inequality implies, for all (x, t) in $Q_{R,T}$

$$(3.15) \quad \begin{aligned} (\psi^2\omega^2)(x, t) &\leq \psi^2(x_1, t_1)\omega^2(x_1, t_1) \leq \psi(x_1, t_1)\omega^2(x_1, t_1) \\ &\leq \frac{\tilde{c}\beta^4 + c(\alpha, \delta, m)}{R^4} + \frac{c(\delta, m)K}{R^2} + \frac{c(\delta)}{T^2} + c(\delta)(|a| + K)^2 + c(\delta)a^2\beta^2. \end{aligned}$$

Note that $\psi(x, t) = 1$ in $Q_{R/2, T/2}$ and $\omega = |\nabla f|^2/(\alpha - f)^2$. Therefore we have

$$(3.16) \quad \frac{|\nabla f|}{\alpha - f} \leq \left(\frac{\tilde{c}\beta^4 + c(\alpha, \delta, m)}{R^4} + \frac{c(\delta, m)K}{R^2} + \frac{c(\delta)}{T^2} + c(\delta)(|a| + K)^2 + c(\delta)a^2\beta^2 \right)^{1/4}.$$

Since $f = \log u$, we get the following estimate for Eq. (2.1)

$$(3.17) \quad \frac{|\nabla u|}{u} \leq \left(\frac{\tilde{c}\beta^4 + c(\alpha, \delta, m)}{R^4} + \frac{c(\delta)}{T^2} + c(\delta)(|a| + K)^2 + c(\delta)a^2\beta^2 \right)^{1/4} (\alpha - \log u).$$

Replacing u by $e^{b/a}u$ gives the desired estimate (1.8). This completes the proof of Theorem 1.2. $\square$

4. PROOF OF COROLLARY 1.4

Proof. The proof is similar to that of Theorem 1.2. We still use the technique of a cut-off function in a local neighborhood of Riemannian manifolds. For $0 < u \leq 1$, we let $f = \log u$. Then $f \leq 0$. Set

$$\omega := |\nabla \log(1 - f)|^2 = \frac{|\nabla f|^2}{(1 - f)^2}.$$

By Lemma 2.1, we have

$$(4.1) \quad \left(L - \frac{\partial}{\partial t} \right) \omega \geq \frac{2f}{1 - f} \langle \nabla f, \nabla \omega \rangle + 2(1 - f)\omega^2 - 2(|a| + K)\omega.$$

We define a smooth cut-off function $\psi = \psi(x, t)$ in the same way as Section 3. Follow all steps in the last section (see also pp. 1050-1051 in [1]), we can easily get

the following inequality

$$(4.2) \quad \begin{aligned} 2(1-f)\psi\omega^2 &\leq (1-f)\psi\omega^2 + \frac{cf^4}{R^4(1-f)^3} + \frac{\psi\omega^2}{2} + \frac{c}{R^4} \\ &\quad + \frac{c(m)}{R^4} + \frac{c(m)K}{R^2} + \frac{c}{T^2} + c(|a| + K)^2, \end{aligned}$$

where we used similar estimates (3.3)-(3.9) with the difference that these estimates do not contain the parameter δ . Using the same method as that in proving Theorem 1.2, for all (x, t) in $Q_{R/2, T/2}$ we can get

$$(4.3) \quad \begin{aligned} \omega^2(x, t) &\leq \frac{c(m)}{R^4} + \frac{c(m)K}{R^2} + \frac{c}{T^2} + c(|a| + K)^2 \\ &\leq \frac{c(m)}{R^4} + \frac{c(m)}{R^2}(|a| + K) + \frac{c}{T^2} + c(|a| + K)^2 \\ &\leq \frac{c(m)}{R^4} + \frac{c}{T^2} + c(|a| + K)^2. \end{aligned}$$

Again, using the same argument in the proof of Theorem 1.2 gives

$$(4.4) \quad \frac{|\nabla f|}{1-f} \leq \frac{c(m)}{R} + \frac{c}{\sqrt{T}} + c\sqrt{K + |a|},$$

where c is a constant depending only on n , $c(m)$ is a constant depending only on n and m .

Since $f = \log u$, we get

$$(4.5) \quad \frac{|\nabla u|}{u} \leq \left(\frac{c(m)}{R} + \frac{c}{\sqrt{T}} + c\sqrt{K + |a|} \right) \cdot \left(1 + \log \frac{1}{u} \right).$$

At last, replacing u by $e^{b/a}u$ above yields (1.9). $\square$

ACKNOWLEDGMENT

The author would like to thank Professor Yu Zheng for his helpful suggestions on this problem, and for his encouragement. He would also like to thank the referees for useful suggestions. This work is partially supported by the NSFC10871069.

REFERENCES

1. Souplet P., Zhang Q S. Sharp gradient estimate and Yau's Liouville theorem for the heat equation on noncompact manifolds. *Bull. London Math. Soc.*, 2006, **38**: 1045-1053.
2. Wu J Y. Elliptic type gradient estimates for a nonlinear parabolic equation on complete manifolds. 2008, preprint.
3. Hamilton R S. Three-manifolds with positive Ricci curvature. *J Diff Geom.*, 1982, **17**: 255-306.
4. Chow B, Lu P, Ni L. Hamilton's Ricci flow, Lectures in Contemporary Mathematics 3, Science Press and American Mathematical Society, 2006.
5. Ma L. Gradient estimates for a simple elliptic equation on non-compact Riemannian manifolds. *J Funct Anal.*, 2006, **241**: 374-382.
6. Yang Y Y. Gradient estimates for a nonlinear parabolic equation on Riemannian manifold. *Proceeding of AMS*, 2008, **136**: 4095-4102.
7. Bakry D. Lhypercontractivité et son utilisation en théorie des semigroupes, 1-114, Lect. Notes in Math., vol. 1581, Springer-Verlag, Berlin/New York, 1994.
8. Bakry D, Emery M. Diffusion hypercontractivites, in Séminaire de Probabilités 19, 1983/1984, 177-206, Lect Notes in Math 1123, Berlin: Springer, 1985.
9. Bakry D, Qian Z M. Volume comparison theorems without Jacobi fields. Current trends in potential theory, 115-122, Theta Ser Adv Math, 4, Theta, Bucharest, 2005.

10. Li X D. Liouville theorems for symmetric diffusion operators on complete Riemannian manifolds. *J Math Pure Appl*, 2005, **84**: 1295-1361.
11. Yau S T. Harmonic functions on complete Riemannian manifolds. *Comm Pure Appl Math*, 1975, **28**: 201-228.
12. Cheng S Y, Yau S T. Differential equations on Riemannian manifolds and their geometric applications. *Comm Pure Appl Math*, 1975, **28**: 333-354.
13. Li P, Yau S T. On the parabolic kernel of the Schrodinger operator. *Acta Math*, 1986, **156**: 153-201.
14. Li J Y. Gradient estimates and Harnack inequalities for nonlinear parabolic and nonlinear elliptic equations on Riemannian manifolds. *J Funct Anal*, 1991, **100**: 233-256.
15. Negrin E. Gradient estimates and a Liouville type theorem for the Schrodinger operator. *J Funct Anal*, 1995, **127**: 198-203.
16. Hamilton R S. A matrix Harnack estimate for the heat equation. *Comm Anal Geom*, 1993, **1**: 113-126.

DEPARTMENT OF MATHEMATICS, EAST CHINA NORMAL UNIVERSITY, SHANGHAI, CHINA 200241
E-mail address: jywu81@yahoo.com